

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 652 709 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
17.03.1999 Bulletin 1999/11

(51) Int. Cl.⁶: A01N 61/00, A01N 43/90,
A01N 43/16

(21) Application number: 93916071.9

(86) International application number:
PCT/GB93/01478

(22) Date of filing: 14.07.1993

(87) International publication number:
WO 94/02022 (03.02.1994 Gazette 1994/04)

(54) IMPROVEMENTS IN OR RELATING TO GERMICIDAL COMPOSITIONS

VERBESSERUNGEN IN, ODER IN VERBINDUNG MIT KEIMTÖTENDEN
ZUSAMMENSETZUNGEN

AMELIORATIONS APPORTEES AUX COMPOSITIONS GERMICIDES

(84) Designated Contracting States:
CH DE ES FR GB IT LI NL SE

(56) References cited:
EP-A- 0 081 462 EP-A- 0 286 524
DE-A- 2 812 261

(30) Priority: 22.07.1992 GB 9215555
30.10.1992 GB 9222813
09.03.1993 GB 9304732

(43) Date of publication of application:
17.05.1995 Bulletin 1995/20

(73) Proprietors:
• UNILEVER PLC
London EC4P 4BQ (GB)
Designated Contracting States:
GB
• UNILEVER N.V.
3013 AL Rotterdam (NL)
Designated Contracting States:
CH DE ES FR IT LI NL SE

- DATABASE CHEMABS CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US AN: CA74(21):108555b K.INADA ET AL. 'Relation between photodynamic activities and photochemical oxidation in xanthene dyes' & SHOKUHIN EISEIGAKU ZASSHI vol. 11, no. 4, 1970 pages 238 - 241
- DATABASE BIOSIS BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US AN: 88:25633; DN: BA85:13358 M.I.ROAT 'THE ANTIVIRAL EFFECTS OF ROSE BENGAL AND FLUORESCIN' & ARCH OPHTHALMOL vol. 105, no. 10, 1987 pages 1415 - 1417
- DATABASE CHEMABS CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US AN: CA72(19):97667v K.INADA ET AL. 'Photodynamic action of food dyes on bacteria' & SHOKUHIN EISEIGAKU ZASSHI vol. 10, no. 5, 1969 pages 344 - 347
- DATABASE CHEMABS CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US AN: CA90(17):134960e S.A.BEZMAN ET AL. 'Photodynamic inactivation of E. coli by rose bengal immobilized on polystyrene beads' & PHOTOCHEM. PHOTOBIOLOG. vol. 28, no. 3, 1978 pages 325 - 329
- S. P. Gorman in: Soc. for Appl. Bacteriol. Techn. Ser., (S. P. Denyer & W. B. Hugo (eds.)), p. 271-295, Oxford, 1991.

(72) Inventors:
• RABONE, Kenneth Leslie
Bebington, Wirral L63 2QP (GB)
• HAQ, Ziya
Gayton, Wirral L60 3RW (GB)

(74) Representative:
Dekker, Enno E. J. et al
UNILEVER PLC,
Patent Division,
Colworth House
Sharnbrook, Bedford MK44 1LQ (GB)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 652 709 B1

- | | |
|--|---|
| <ul style="list-style-type: none">• M. R. W. Brown & P. Gilbert, J. Appl. Bacteriol. Symp. Suppl., Vol. 74, 87S - 97S (1993).• D. S. Dhaliwal et al., Lett. Appl. Microbiol., Vol. 15, 217 - 221 (1992).• J. T. Holah et al., Lett. Appl. Microbiol., Vol. 11, 255 - 259 (1990). | <ul style="list-style-type: none">• J. T. Patterson, J. Food Technol., Vol. 6, 63 - 72 (1971).• P. Maris, Sci. Aliments., Vol. 12 (4), 721 - 728 (1992). |
|--|---|
-

Description

Field of the Invention

- 5 [0001] This invention relates to germicidal compositions, particularly for use on surfaces, ie compositions capable of destroying or inactivating micro-organisms, particularly surface-bound micro-organisms.

Summary of the Invention

- 10 [0002] In its broadest aspect the present invention provides a method of killing bacteria in biofilms on a surface, comprising applying to the surface a composition comprising a dyestuff which is capable of photo-dynamic inactivation of micro-organisms.

- [0003] It is preferred to use a dyestuff that generates singlet oxygen on exposure to light. On absorption of light energy a dye molecule is converted to a more energetic or excited state (S_1) from its electronic ground state (S_0). (Electron spins are paired and so these are singlet states in the language of spectroscopy, having a single energy level in a magnetic field.)

[0004] The excited state is short lived and can lose energy and return to the ground state in a number of ways: by emission of a quantum of light as fluorescence; by internal conversion as the energy is degraded to heat, by collision with a molecule of a different substance (fluorescence quenching).

- 20 [0005] The short lived singlet state may also undergo a process called intersystem crossing to a longer-lived excited state, the triplet state. (The state is termed "triplet" because the electron in the higher energy level is no longer spin-paired with the electron in the lower level and the excited state has three energy levels in a magnetic field.)

- [0006] Interaction of the excited triplet state with ground state molecular oxygen (which exists in the triplet state normally) regenerates the dye ground state as energy is transferred to the oxygen which is promoted to the electronically excited singlet state. This means that, under ideal circumstances, a single photosensitiser molecule can generate many times its own concentration of singlet oxygen.

- [0007] Singlet oxygen is highly reactive and photosensitised oxidation proceeding via this route is known as Type II photo-oxidation. Type II photo-oxidation is independent of the photosensitiser used to generate the singlet oxygen. An important feature of the sensitiser is that it should have a high quantum yield of triplet formation (that is, ideally, a triplet state should be produced for each photon absorbed). Intersystem crossing to the triplet state is facilitated by the presence of heavy atoms in the molecule.

[0008] Photo-oxidation of any vital component of an organism may result in cell death (protein, polypeptide, amino-acids, lipids with allylic hydrogens, tocopherols, sugars and cellulose).

- [0009] Currently preferred dyestuffs include Rose Bengal (Acid Red 94, Colour Index No. 45440), Erythrosin B (Acid Red 51, Colour Index No. 45430), and phthalocyanin sulphonates such as aluminium phthalocyanin sulphonate (APS) and zinc phthalocyanin sulphonate (ZPS). Rose Bengal and Erythrosin B are known food colourants (Rose Bengal is Food Colour Red No 105 and Erythrosin B is Food Colour Red No 14), and Erythrosin B is on an EEC list of colouring agents allowed for use in cosmetic products, so these two dyes are well suited for use in compositions intended for domestic use. Mixtures of dyes can be used, and in some cases it may be desirable to include in a mixture a dye that will remain visible at the end of the photodynamic process.

[0010] The concentration of dyestuff in the composition is not critical and will typically be up to 100ppm, with good results having been obtained with concentrations in the range 10ppm to 20ppm. Lower concentrations, down to 1ppm should also give reasonable results.

- [0011] Singlet oxygen has a short lifetime and therefore a short pathlength for diffusion, so to be effective a photosensitising dye generating singlet oxygen must be close to the target substrate. Preferred dyestuffs are therefore substantive to (ie capable of binding to) micro-organisms, typically by binding to cellular protein on the organism surface or other cellular components eg cellular fats.

- [0012] The preferred dyes mentioned above generate singlet oxygen on exposure to light and are substantive to protein and so capable of binding to micro-organisms via cellular protein. In this way, targetted killing of organisms and hence germicidal action is possible.

- [0013] It is also preferred to use a dyestuff that is bleached by exposure to light. By use of a photo-bleaching dyestuff that is substantive to micro-organisms it can be possible for a visible indication of the presence of micro-organisms to be provided. As the dyestuff bleaches, the photo-dynamic action proceeds causing the death of, or otherwise inactivating, micro-organisms. In the presence of low levels of light, both bleaching and the photo-dynamic activity are believed to proceed more slowly, whereas at higher light intensities both processes occur more quickly. Thus, depending on the relative rates of bleaching and photo-dynamic activity, the presence of visible dyestuff indicates to the user that the photo-dynamic inactivation of any micro-organisms present is incomplete.

[0014] The photo-dynamic inactivation of micro-organisms in suspension by dyes such as Rose Bengal is known.

See, for example, Journal of Applied Bacteriology 1985, [58] pages 391-400, Photochemistry and Photobiology 1988, [48] pages 607-612 (Neckers *et al*), and Shokuhin Eiseigaku Zasshi 1962, 10(5), pages 344-347. See also EP-A-81462 which describes novel water-soluble zinc and aluminium phthalocyanines and their use as photoactivators in microbial compositions. Example 9 concerns use of the compounds on a suspension of bacteria placed on a test surface. However, it has now been found that suitable dyestuffs are capable of photo-dynamic inactivation of micro-organisms on surfaces, and this is the basis of the present invention. It is well-known that micro-organisms are much more susceptible to biocides in their planktonic or suspended state: they are much more difficult to inactivate when attached to surfaces, which is their usual or preferred state. Micro-organisms will normally be on surfaces in the form of "biofilms", that is, embedded in a matrix of extracellular material. This extracellular material may sometimes be referred to as "adhesin" in the literature. It is therefore not obvious that a process which acts on micro-organisms in their planktonic state would act on surface-bound organisms without modification being required. Surface-bound micro-organisms represent an important and substantial source of contamination in domestic, institutional and industrial environments, and the present invention can enable targetted germicidal action on such micro-organisms.

[0015] The method of the invention is particularly suitable for use in killing micro-organisms on hard domestic and industrial surfaces such as glass, plastics, ceramic and metal surfaces. In particular, the method is effective on surfaces which may harbour soils having the potential for bacteriological contamination in surface imperfections, joints and other relatively confined regions.

[0016] The composition is preferably acidic, eg having a pH in the range of 3 to 5, eg a pH of about 4, as acidic compositions are found to have substantially enhanced effectiveness against Gram-negative (G-) micro-organisms as compared with neutral compositions. The effectiveness against Gram-positive (G+) micro-organisms seems not to be significantly affected by pH. The compositions are conveniently made acidic by use of relatively mild organic acid, such as acetic acid.

[0017] Neckers *et al* (above) review conflicting evidence that penetration of the dye Rose Bengal itself through the cell wall is essential for inactivation. They suggest that their own results support the dye penetration hypothesis primarily on the basis of differential inactivation of G+ and G- species. The envelope of G- bacteria has an additional outer membrane composed substantially of lipopolysaccharide which Necker *et al* regard as serving as a barrier to potentially toxic substances. An alternative interpretation could be that the amount of protein exposed in the cell walls is very different between G+ and G- species, being greater for G+ than G-, with a binding affinity for the dye which varies with pH. The concentration of dye bound to the cell wall would therefore be a function of pH. This interpretation would account for our observations of the variation in kill rate with pH (but not the apparent resistance of the G+ species *B. subtilis* and *B. megaterium*).

[0018] There is no difference between the endospore-forming species *B. subtilis* and the nonspore-forming *S. aureus* in their susceptibilities to photodynamic action by Rose Bengal in the absence of endospores. The apparent differences shown in our studies are presumably due to the presence of endospores both for *B. subtilis* and *B. megaterium* which are clearly more resistant to singlet oxygen than the bacteria themselves. The spores survive exposure to Rose Bengal and light and subsequently germinate to give countable colonies. Spores are known to be difficult to stain and presumably have no affinity for Rose Bengal under any of the conditions used. There appears to be little documentation on the toxicity of singlet oxygen to spores in the literature. From studies on the possibly less resistant conidia of *Neurospora crassa* (Photochem. Photobiol., 33, 349 (1981)), singlet oxygen does have potential in this respect.

[0019] The composition may optionally include other ingredients such as one or more surfactants (for cleaning purposes) and/or one or more solvents.

[0020] The surfactant is preferably alkoxylated, more preferably ethoxylated, eg being in the form of ethoxylated alcohols. The alcohol preferably has between 4 and 15 carbon atoms, is of straight or branched chain configuration, and has an HLB value (hydrophilic lipophilic balance) in the range 10 to 14, eg 12.

[0021] A wide range of suitable surfactants are commercially available, one such material being the surfactant available under the trade name Imbentin 91-35, from Kolb, which is a nonionic C9-11 alcohol ethoxylate, having an average of 5 moles of ethylene oxide per mole of alcohol.

[0022] Primary ethoxy sulphates may also be used.

[0023] Mixtures of surfactants may be used if desired.

[0024] The surfactant is preferably non-ionic or anionic, or a mixture of both types.

[0025] Preferred anionic surfactants for this purpose include primary alkyl sulphates (PAS), preferably sodium dodecyl sulphate (SDS). Commercial mixtures containing a substantial proportion of dodecyl sulphate (eg Empicol LX) are especially preferred. Dodecyl sulphate is a known protein denaturant, is good for cleaning protein off surfaces, and is biocidal.

[0026] The composition is preferably substantially free of cationic surfactant, but may include a minor amount of cationic germicide.

[0027] Surfactant preferably constitutes an amount in the range 0.05 to 2.5% by weight of the total weight of the composition, typically 0.5% to 1.5% by weight, eg 0.7% by weight nonionic surfactant with an optional amount of up to 0.2%

by weight of anionic surfactant.

[0028] The solvent is preferably polar and is preferably a straight or branched chain C2 to C5 alcohol such as ethanol, butanol, isopropanol (propan-2-ol) (IPA), N-butoxy propan-2-ol (propylene glycol n-butyl ether), 2-butoxy ethanol (ethylene glycol monobutyl ether). IPA is the currently preferred solvent.

[0029] Dihydric alcohol such as ethylene glycol, and water miscible ethers such as dimethoxyethane may also be used.

[0030] Mixtures of solvents can be used if appropriate, eg mixtures of ethanol and N-butoxy propan-2-ol.

[0031] Solvent is preferably present in an amount in the range 2 to 20% by weight of the total weight of the composition.

[0032] At least some of these solvents, eg ethanol, weaken the cell walls of micro-organisms, making them more permeable and so more susceptible to penetration by singlet oxygen. This has the effect of enhancing the micro-organism-killing effect of the dyestuff.

[0033] It is found that inclusion of surfactant can reduce the photo-dynamic effect of dyestuffs (possibly by solubilising the dyestuff and preventing adsorption on the cell wall), and inclusion of solvent can also reduce the photo-dynamic effect of dyestuffs (possibly by competing for singlet oxygen). However, it is found that with three component formulations, comprising dyestuff, surfactant and solvent, the reduction in photo-dynamic effect of the dyestuff is less than would be expected from the combined effects of surfactant and solvent. The surfactant and solvent together thus have a synergistic effect, the result of which is to lower the reduction in the photo-dynamic effect of the dyestuff.

[0034] In a preferred aspect, the present invention thus provides a method of cleaning a surface and killing bacteria in biofilms on the surface, comprising applying to the surface a composition a dyestuff which is capable of photo-dynamic inactivation of micro-organisms, a surfactant and a solvent.

[0035] The composition may include a number of optional ingredients including the following:

1. Detergent boosters, preferably metal chelating agents such as ethylene diamine tetra acetic acid (EDTA). Metal chelating agents (including EDTA) have also been claimed to permeabilise cell walls, thus making organisms more susceptible to biocides.

2. Electrolyte such as a buffer or salt, eg Na_2SO_4 , which acts to assist binding of dye to protein by promoting movement of dye from the aqueous phase to the protein salt. Electrolyte is commonly present in various dye formulations as commercially available, although additional electrolyte can be added if required. Total electrolyte can be added if required. Total electrolyte content of the composition would typically be in the range 0 to 1% by weight, preferably about 0.1%.

3. Perfumes.

4. Thickeners.

[0036] The composition is in the form of an isotropic, single phase composition and is of particular use as a germicide (possibly also with a cleaning effect) on hard surfaces, finding application in a wide range of contexts, including domestic applications, eg kitchen and bathroom surfaces including toilet bowls, in institutions such as schools, hospitals etc, and in commercial premises such as factories, offices, hotels etc.

[0037] For domestic use at least, the composition is preferably formulated as a product intended for application by spraying and is conveniently packaged in a suitable container, eg having a hand operated trigger spray or an aerosol propellant dispenser. The container is preferably light-opaque.

[0038] The methods of the invention involve applying to a surface to be treated in any convenient manner, eg by spraying from a suitable dispenser, wiping on with a carrier such as a cloth or sponge, or pouring from a container etc. This might in some cases, particularly in industrial cleaning, be followed by exposure to a light source, eg a white light source such as a quartz halogen lamp or fluorescent "daylight" source. (The process would be an alternative to using dangerous germicidal radiation, for example from a low pressure mercury discharge lamp emitting resonance radiation at 254 nm. Such radiation is harmful to the unprotected eye.) This would generally be followed by a rinsing step, if required, eg by wiping with a carrier, application of a stream of running water etc.

[0039] The invention will be further described, by way of illustration, in the following Examples and by reference to the accompanying Figures in which:

Figure 1 shows two bar charts of log (reduction) values illustrating the lethal effect of Rose Bengal and light on various micro-organisms in suspension at pH4 and pH7, with Figure 1a showing results for Gram-positive micro-organisms, and Figure 1b showing results for Gram-negative organisms and yeasts;

Figure 2 is a pair of graphs of log (reduction) versus pH showing the biocidal effect of Rose Bengal and light on *S. aureus* and *E. coli* as a function of pH, with Figure 2a showing results after 20 minutes exposure to light and Figure 2b showing results after 60 minutes exposure;

Figure 3 is a pair of graphs similar to Figure 2 obtained using Erythrosin B in place of Rose Bengal;

Figure 4 shows two bar charts of log (reduction) illustrating the photohygiene effect of various combinations of Rose Bengal (RB), Imbentin C91-35 (AE), isopropanol (IPA) and Empicol LX (PAS), with Figure 4a showing results obtained without exposure to light and Figure 4b showing results obtained with exposure to light;

Figure 5 is a graph of adsorption of Rose Bengal by *E. coli*, with amount adsorbed (nanomoles) versus equilibrium concentration (micromoles/l), with results at pH 4 shown by squares and results at pH 7 shown by crosses;

Figure 6 is a pair of graphs similar to Figure 5 showing the effect of electrolyte, with Figure 6a showing results at pH 4 and Figure 6b showing results at pH 7, with results without additive shown by squares, results with sodium sulphate (1%) shown by crosses and results with sodium sulphate (5%) shown by double crosses;

Figure 7 is a pair of graphs similar to Figure 5 showing the effect of surfactant, with Figure 7a showing results at pH 4 and Figure 7b showing results at pH 7, with results without additive shown by squares, results with non-ionic surfactant (0.7%) (NI) shown by crosses and results with PAS (0.7%) shown by double crosses;

Figure 8 is a graph similar to Figure 5 showing the effect of solvent at pH 4, with results without additive shown by small squares, those for 10% IPA shown by crosses, those for 0.7% Imbentin shown by double crosses and those for 10% IPA and 0.7% Imbentin shown by large squares;

Figure 9 is a graph of log (reduction) versus exposure time (minutes) showing the effect of pH on the rate of kill of *E. coli* by Rose Bengal, with results at pH 4 shown by squares and results at pH 7 shown by crosses; and

Figure 10 is a graph similar to Figure 9 showing the effect of electrolyte at pH 7 on the rate of kill of *E. coli* by Rose Bengal, with results without electrolyte shown by squares and results with Na sulphate shown by crosses.

Examples

EXPERIMENTAL PROCEDURES

Preparation of Inocula

[0040] The following micro-organisms (generally either from the National Collection of Type Cultures (NCTC) or the American Type Culture Collection (ATCC)) were used in the experiments described:

Bacteria:

[0041]

<i>Staphylococcus aureus</i>	NCTC 6538 (Gram positive)
<i>Escherichia coli</i>	NCTC 8196 (Gram negative)
<i>Pseudomonas aeruginosa</i>	NCTC 5940 (Gram negative)
<i>Enterobacter sp.</i>	NCTC 3281 (Gram negative)
<i>Klebsiella sp.</i>	ATCC 11677 (Gram negative)
<i>Bacillus subtilis</i>	NCTC 6432 (Gram positive)
<i>Bacillus megaterium</i>	NCTC 7581 (Gram positive)

Yeast:

[0042]

5 Candida albicans

[0043] Organisms were grown up by overnight incubation in nutrient broth at 37°C for bacteria (28°C for *Ps. aeruginosa*) or SABS broth (SABS is Sabourand Dextrose Agar, with liquid medium in the case of SABS broth, from Oxoid Ltd) at 28°C for yeast. Cultures were isolated by vacuum filtration using a 0.45µm Millipore filter and washed with quarter-strength Ringers solution before resuspension in Ringers solution (10ml). The organisms in suspension were enumerated by serial dilution and plating with nutrient agar (bacteria) or SABS agar (yeast) and the total viable count (TVC) expressed as the decadic logarithm of the number of colony-forming units (cfu) per ml.

[0044] Experiments were carried out at pH 7 unless otherwise specified.

15 1) AGAR DIFFUSION DISC METHOD

[0045] Aqueous solutions containing 100 ppm of dye were prepared. Aliquots (10ml) of each dye solution were sterilized in glass universal screw cap vials. Antibiotic assay discs (13mm from BDH) were also sterilized. All organisms were grown overnight in nutrient or SABS broth (10ml).

20 [0046] For each micro-organism, two nutrient agar plates (SABS agar for the yeast) were seeded with the overnight culture (10ul) to give confluent growth over the whole plate. Using aseptic techniques, an antibiotic disc was dipped into the first dye solution and placed on the surface of a seeded agar plate. This was repeated with two other dye solutions to give three discs on duplicate plates.

25 [0047] One of each pair of duplicate plates was immediately placed in an incubator at the appropriate temperature with minimal exposure to light. The remaining plates were placed on top of a light box for 3 hours. Illumination from the light box was diffused white light of mean intensity 4000 lux from fluorescent "daylight" tubes (2 x 15 watt, Exal X-ray Accessories Ltd, Hemel Hempstead). Light intensities were measured at the surface of the diffuser using a Megatron DA10 light meter. After exposure the plates were incubated overnight and then examined for zones of inhibition around the disc.

30 [0048] Using this approach, the following results were obtained:

Example 1 (Disc Method)

35 [0049] Results on agar for aqueous dye solutions of Rose Bengal, Erythrosin B and aluminium phthalocyanin sulphate (APS) (100 ppm) at pH 7 after exposure to light for 180 minutes as described above are summarised in Table 1. In the Table, the results are expressed as the difference (in millimetres) between the radius of the clear zone of inhibition (the area of no bacterial growth on the spread agar plate) and the radius of the disc. Thus, the higher the value, the greater the bacterial kill.

40 [0050] The agar diffusion disc method ranked Rose Bengal as more effective than the structurally similar Erythrosin B. It is tempting to ascribe this ranking to a difference in the quantum yield for singlet oxygen formation. In methanol, the quantum yield for singlet oxygen formation is 0.76 for Rose Bengal compared to 0.6 for Erythrosin B. However, a number of other factors might also be expected to contribute to the observed differences such as the rate of dye diffusion or differences in dye binding to the agar gel or disc material.

45 [0051] A particular feature of the results from the disc method is the clear distinction in the sensitivity of Gram-positive (G+) and Gram-negative (G-) organisms to photodynamic action, at least at pH 7. Possible reasons for the relative resistance of G- organisms are discussed elsewhere.

2) SURFACE TEST

50 Test Solutions

[0052] Rose Bengal (20ppm) in pH 4 buffer and:

1. No further addition
- 55 2. Ethanol (10 % v/v) and Imbentin C91-35 (0.7%).
3. Propan-2-ol (10% v/v) and Imbentin C91-35 (0.7%).

[0053] Sodium hypochlorite solution (0.125 %) was used as a positive control.

Estimation of Number of Organisms Adherent to Base of Petri Dish

[0054] Micro-organism, eg *S. aureus*, suspension (0.5ml) was added to aliquots (100 ml) of quarter-strength Ringers solution and the average cfu per ml determined (TVC). Aliquots (20ml) of these solutions were pipetted into sterile petri dishes and left at room temperature for 5 hours. The inoculum was then removed by pipette into a sterile bottle and the average cfu per ml remaining in suspension estimated. The number of organisms (as cfu) per square cm adherent to the Petri dish was calculated from the difference in the solution concentrations.

Test Method

[0055] Bacterial suspensions in quarter-strength Ringers (20mls) were pipetted into sterile plastic Petri dishes which were left at ambient temperature for 5 hours. The inoculum was then removed, the dishes washed once with Ringers solution with gentle swirling by hand, and the solution poured off. Pairs of bacterially contaminated plates were treated with the test solutions. An aliquot (20ml) of test solution was poured into one dish and the solution decanted after 30 seconds. The dish was rinsed with pH 4 buffer and placed on the light box for 20 mins. In a separate experiment, the solution was exposed in the second dish on the light box for 20 minutes before being decanted and the dish rinsed with pH 4 buffer. Both experiments were duplicated.

[0056] After light exposure, one of the duplicate plates was overlaid with Tryptone Soya agar containing 1% glucose and 0.015% Neutral Red cooled to about 50°C. The other duplicate plate was stained with 0.01% Acridine Orange for 30 seconds, rinsed and examined microscopically (Nikon "Optiphot" microscope equipped with a 100x apochromat oil-immersion objective, 10x eyepiece and epifluorescence attachment with a B2-A combined filter/dichroic mirror block and super high pressure mercury lamp). The overlaid agar plates were incubated at 37°C for 48 hours, by which time colonies had grown out of the adherent bacteria which had not been killed. The colonies took up the Neutral Red and could be seen and counted under the agar on the surface of the dish. (A M R MacKenzie and R L Rivera-Calderon, Agar Overlay Method to Measure Adherence of *Staphylococcus epidermidis* to Four Plastic Surfaces, Applied and Environmental Microbiology, 50, 1322 (1985).)

[0057] Plates stained with Acridine orange were examined and photographed. The number of stained bacteria were counted in a field of view (clumps counted as one) which had previously been estimated by photographing a micrometer scale.

Example 2 (Surface Test)

[0058] Control experiment surface tests using both direct epifluorescence microscopy as described above and suspension depletion gave similar values for the number of bacteria (*S. aureus*) that could be attached to the surface of a plastic dish (of the order of one million per square centimetre).

[0059] Contracting the surface with the positive control (sodium hypochlorite solution, 30 seconds) reduced the number of viable bacteria to zero. The results for photo-dynamically inactivated bacteria are summarised in Table 2 as the decadic logarithm of the reduction (log (reduction)) of viable bacteria before and after exposure, ie log (start count) - log (final count).

Example 3

[0060] Formulation: Rose Bengal (20 ppm), nonionic surfactant (Imbentin C91-35, 0.7%), propan-2-ol (10 %) (pH 4).

[0061] Experimental procedure was as previously described except that surface attachment of the bacteria required them to be in the exponential growth phase. This was achieved by incubation in nutrient broth (for bacteria) or SABS broth (for yeast) in the plastic Petri-dish for 3 hours. In Example 2, a suspension of non-growing bacteria in Ringers' solution was simply allowed to stand for 5 hours. This worked well for *S. aureus* but not other bacteria. The results are shown in Table 3.

[0062] Previous experience using direct epifluorescence microscopy suggests that where confluent growth was obtained, this would be equivalent to an initial surface density of viable microbes of the order of 1 million/sq. cm. The surface area of the Petri-dishes used was about 57 sq.cm. so that the treatment in all cases has reduced the level of surface contamination by at least 5 orders of magnitude (log 5).

[0063] Surface tests are more difficult to perform than suspension tests, and for this reason most experimental work has been performed on bacteria in suspension. Although the following Examples are not within the scope of the invention, they are included to demonstrate the effect of varying conditions.

3) SUSPENSION TEST

Preparation of Solutions

- 5 [0064] Stock solutions of the following were prepared by weighing and sterilized (except those containing solvent):
 [0065] Rose Bengal (0.2 percent) in propan-2-ol (95 percent) Nonionic surfactant (Imbentin C91-35, 14 percent) (sometimes referred to by the abbreviation AE, for alcohol ethoxylate)
 Anionic surfactant (Empicol LX, 14 percent) (sometimes referred to as PAS)
 pH 4 buffer (citric acid (0.1 M, 307 ml) + dibasic sodium phosphate (0.2 M, 193 ml)
 10 pH 7 buffer (sodium dihydrogen orthophosphate (0.4M, 468 ml) + disodium hydrogen orthophosphate dodecahydrate, (0.4M, 732ml)
 Buffers pH 5, 6, 8, 9 were prepared as indicated in the CRC Handbook of Chemistry and Physics, 8-36, 73rd Ed., CRC Press (1992-1993)
 [0066] Final concentrations in the test solutions were typically:
- 15 Rose Bengal - 20ppm
 Ethanol - 10.0% (v/v)
 Surfactant - 0.7% (w/v)
- 20 [0067] In some examples, different concentrations were used as specified.

Test Method

- 25 [0068] Test solutions were made up in sterile plastic petri dishes to a depth of 5mm (30mls). A suspension of micro-organism (0.3ml) was added to each solution and gently mixed in. If Rose Bengal was to be included in the test solution it was added last to minimise light exposure. Solutions were either exposed on a light box, placed in the dark (conditions of reduced light exposure) or left on the bench. The average intensity at the surface of the light box diffuser was 4000 lux measured with a Megatron DA 10 light meter (from Megatron Ltd). After specified exposure times, surviving bacteria were enumerated as colony-forming units (cfu/ml) following incubation after serial dilution and plating onto agar.
 30 The decadic logarithm of the number of bacteria remaining (as colony-forming units per ml) was determined and compared to the number before exposure as log (start count) - log (final count). The higher the value, the greater the bacterial kill.
 [0069] Suspension tests have been carried out against a variety of organisms under a variety of conditions to optimise photo-dynamic action against a range of micro-organisms, including Gram-negative organisms and yeast. Salient results are summarised in the associated Tables. In these Tables results are expressed as the decadic logarithm of the
 35 ratio of the initial number of colony-forming units per ml to the number remaining after exposure, the log (reduction). Using this notation a value of zero means no change in the number of organisms following exposure to the conditions. The notation "+" preceding a log reduction figure indicates that no micro-organism growth could be observed (ie total kill).

40 Example 4

- [0070] Suspension tests were carried out as described above to show the lethal effect of Rose Bengal and light on micro-organisms in suspension; particularly the synergy of low pH and ethanol solvent. Tests were carried using Rose Bengal (20ppm) alone or with ethanol (10% (v/v), with light exposure of 20 minutes (light box). The results expressed
 45 as log (reduction) values are given in Table 4.
 [0071] Longer exposure times (of 60 and 100 minutes) improve performance for Gram negative organisms, particularly at pH 7.
 [0072] It will be seen that performance for Gram negative organisms is much improved at pH 4 as compared with pH 7.

50

Example 5

- [0073] Suspension tests were carried out as described above at pH 4 and pH 7 to show the lethal effect of Rose Bengal (20ppm) and light (20 minutes exposure on a light box) of various Gram-positive and Gram-negative micro-organisms and yeast in suspension, and the results expressed as log (reduction) are shown graphically in Figures 1a and 1b, with Figure 1a giving results for Gram-positive micro-organisms and Figure 1b giving results for Gram-negative micro-organisms and yeast.

Example 6

[0074] Suspension tests were carried out as described above at a range of different pHs to show the biocidal effect of Rose Bengal (20ppm) and light on S. aureus (G+) and E. coli (G-) as a function of pH, and the results expressed as log (reduction) are shown graphically in Figures 2a (exposure time 20 minutes) and 2b (exposure time 60 minutes). In the Figures, crosses show results for control (G-), double crosses show results for E. coli, inverted triangles show results for control (G+) and triangles show results for S. aureus.

[0075] Similar suspension tests were carried out to show the biocidal effect of Erythrosin B and light on S. aureus and E. coli as a function of pH, and the results are shown graphically in Figures 3a and 3b, which are otherwise identical to Figures 2a and 2b. These show that Erythrosin B performs similarly to Rose Bengal in terms of its photobiocidal profile with pH.

Example 7

[0076] Further suspension tests were carried out as described above at pH 7 using the following solutions:

Nonionic surfactant Imbentin C91-35 (0.7%)
 Imbentin C91-35 (0.7%) with Rose Bengal (100 ppm)
 Anionic surfactant Empicol LX (0.7%)
 Empicol LX with Rose Bengal (100 ppm)
 Imbentin C91-35, Empicol LX (both 0.7%) and Rose Bengal (100 ppm).

[0077] The results are set out below in Table 5. In the Table the term "dark" indicates conditions of reduced light exposure rather than total darkness because of the practical difficulties of avoiding some light exposure. In the Table the term "light" indicates results that are the average of several experiments carried out over a range of times (20 mins, 1 hour, 3 hours) in a statistically designed experiment.

[0078] Results obtained in similar manner for E. coli at pH 7 are shown graphically in the bar charts of Figure 4. In this Figure PAS is used as an abbreviation for Empicol LX, IPA is used as an abbreviation for isopropanol and AE as an abbreviation for Imbentin C91-35. This figure represents averaged data for AE 0.7%, PAS 0.7%, IPA 10%.

Example 8

[0079] Further suspension tests were performed at pH 4 for E. coli using one or more of the following: Rose Bengal 40ppm, surfactant 0.7% (Imbentin C91-35 or Empicol LX), IPA (isopropanol) 10%. Results are given in Table 6.

[0080] The following examples concern suspension tests carried out generally as described above, but at pH 4. Rose Bengal when used was present at a concentration of 20ppm, although in some cases control solutions without Rose Bengal were exposed to light and the results for these are given in the column headed "No Rose Bengal".

[0081] Examples 9, 10, 11 and 12 used the Gram positive organism S. aureus, and Examples 13 and 14 the Gram negative organism E. coli. Other reagents used are indicated in the examples. In all these Examples, samples were exposed for 20 minutes on a light box. The average intensity at the surface of the diffuser was 4000 lux measured with a Megatron DA10 light meter (from Megatron Ltd).

Example 9

[0082] Suspension tests were carried out using Rose Bengal, ethanol and Imbentin C91-35, with S. aureus. The decadic logarithm of the starting concentration, log (start), of S. aureus was 6.8. Results are given in Table 7.

Example 10

[0083] Suspension tests were carried out using Rose Bengal, Dowanol PnB and Imbentin C91-35, with S. aureus. The log (start) was 6.9. Results are given in Table 8.

[0084] This example shows that Dowanol PnB has certain biocidal properties.

Example 11

[0085] Suspension tests were carried out using Rose Bengal, ethylene glycol and Imbentin C91-35, with S. aureus. The log (start) was 6.8. Results are given in Table 9.

Example 12

[0086] suspension tests were carried out using Rose Bengal, IPA and Lialet 111, with S. aureus. Lialet 111 is the trade name of an ether sulphate formulation commercially available from Enichem, having an average chain length 11 with an average degree of ethoxylation of 3. The log (start) was 6.7. Results are given in Table 10.

Example 13

[0087] Suspension tests were carried out using Rose Bengal, propan-2-ol and Imbentin C91-35, with E. coli. The log (start) was 6.8. Results are given in Table 11.

Example 14

[0088] Suspension tests were carried out using Rose Bengal, ethanol and Imbentin C91-35, with E. coli. The log (start) was 7.1. Results are given in Table 12.

Example 15

[0089] Rose Bengal adsorption was determined from the depletion in solution concentration. Concentrations were obtained spectroscopically from absorbances measured at the wavelength of maximum absorbance (ca. 549 nm) using a WPA Linton S110 spectrophotometer on supernatant liquors freed from microbes by centrifugation.

[0090] The results are shown in Figures 5 to 8.

[0091] The results show that the photobiocidal effect is dependent on dye adsorption. Dye adsorption is:

- a) increased by low pH (Figure 4);
- b) increased by neutral electrolyte (which also increases the photobiocidal effect at neutral pH on E. coli) (Figure 6);
- c) decreased by surfactant (Figure 7);
- d) increased by propan-2-ol (Figure 8).

[0092] Calculations show the amount adsorbed onto E. coli is of the right order of magnitude for monolayer coverage, given that dyes are known to aggregate and the calculated surface area of E. coli must be an underestimate (no account taken of fimbriae/pili). Brief details of the calculations are given below. The molecular dimensions of Rose Bengal were taken from a scale model (Catalin Ltd., London).

Surface area of E. Coli 1E7 sq.nm

Area of Rose Bengal 2 sq.nm (flat)
0.5 sq.nm (side)

Monolayer coverage 5E6 (flat) or 20E6 (side)

Measured adsorption 2 - 8 E-16 Moles/bacteria

Number of molecules 120 - 480 E6 per bacterium

Example 16

[0093] Suspension tests were carried out as described above to determine the effect on the rate of kill of E. coli by Rose Bengal (20 ppm) of pH and addition of electrolyte (Na sulphate, 5%) at pH 7, and the results are shown graphically in Figures 9 and 10, respectively.

Table 1

(Example 1) Photobiocidal Effect of Disclosing Agents Zone of Inhibition Around Disc				
Organism	Gram Type	Rose Bengal	Erythrosin B	APS
<u>S. aureus</u>	+	4	1	2
<u>B. subtilis</u>	+	3	1	1
<u>B. megaterium</u>	+	3	1	1.5
<u>E. coli</u>	-	0	0	0
<u>K. pneumoniae</u>	-	1.5	0	3
<u>Ps. aeruginosa</u>	-	0	0	0
<u>Enterobacter sp.</u>	-	0	0	0
<u>C. albicans</u>		0	0	0

Table 2

(Example 2) The Lethal Effect of Rose Bengal and Light on Staphylococcus Aureus Attached to a Plastic Surface	
Solution	Log (Ratio)
Rose Bengal	4.7
Rose Bengal + Imbentin C91-35 + Ethanol	6.0

Table 3

(Example 3)			
Organism	Experiment Number	Results from Agar	Overlay Technique
		Control	After Exposure
<u>S. aureus</u>	1	Confluent growth	No growth
	2	Confluent growth	No growth
<u>E. coli</u>	1	Much growth	45 cfu
	2	Confluent growth	No growth
<u>K. pneumoniae</u>	1	Confluent growth	No growth
	2	Confluent growth	No growth
<u>P. aeruginosa</u>	1	Much growth	5 cfu
	2	Confluent growth	No growth
<u>C. albicans</u>	1	Confluent growth	5 cfu
	2	Confluent growth	4 cfu

Table 4

(Example 4)					
Organism	Gram Type	Conditions		of Exposure	
		No Solvent		With	Ethanol
		pH 7	pH4	pH 7	pH4
<u>S. aureus</u>	+	7.0	7.1	7.1	6.9
<u>B. subtilis</u>	+	0.6	2.1	3.1	0.8
<u>B. megaterium</u>	+	1.3	0.3	1.1	0.3
<u>E. coli</u>	-	0.2	5.3	0.1	6.9
<u>K. pneumoniae</u>	-	0.9	5.6	0	7.0
<u>Ps. aeruginosa</u>	-	0	7.0	0	6.9
<u>Enterobacter sp.</u>	-	1.1	7.3	0	7.4
<u>C. albicans</u>	-	0	5.7	0	5.7
Controls (No Rose Bengal)					
<u>E. coli</u>	-		0.1		0.8

Table 5 (Example 7)

ORGANISM	IMBENTIN		IMBENTIN/RB		EMPICOL		EMPICOL/RB	
	light	dark	light	dark	light	dark	light	dark
<u>St. aureus</u>	4.5	4.0	8.0	3.5	4.5	4.5	5.5	5.0
<u>E. coli</u>	9.0	9.5	10.5	10.0	5.5	5.0	6.5	6.0
<u>Enteroc.</u>	0	0	0	0	0	0	0	0
<u>Klebsiella</u>	7.0	6.0	6.0	7.0	6.0	6.0	6.0	6.0
<u>Ps. aerug.</u>	0	0	0	0	0	0	0	0
<u>C. albicans</u>	0	0	1.0	1.0	2.5	1.5	3.0	3.0

ORGANISM	IMBENTIN/EMPICOL		IMBENTIN/EMPICOL/RB		ROSE BENGAL	
	light	dark	light	dark	light	dark
<u>St. aureus</u>	3.0	4.0	2.0	0	4.0	0
<u>E. coli</u>	4.0	4.0	4.0	4.5	0	0
<u>Enteroc.</u>	0	0	0	0	0	0
<u>Klebsiella</u>	3.5	3.5	4.0	2.5	0	0
<u>Ps. aerug.</u>	0	0	0	0	0	0
<u>C. albicans</u>	0	0	0	0	0	0

Table 6

(Example 8)						
	EXPOSURE TIME					
	1 HOUR		2 HOURS		3 HOURS	
	light	dark	light	dark	light	dark
Rose Bengal	3.8	0.5	7.2	0.4	7.2	0.7
RB/Imbentin C91-35	0.2	0.2	1.0	0.3	3.8	0.5
RB/Empicol LX	1.7	0.4	3.9	0.7	7.2	0.7
RB/IPA	5.4	3.6	4.7	5.0	7.2	7.2
Imbentin C91-35	0.1		0.3		0.9	

Table 6 (continued)

(Example 8)						
	EXPOSURE TIME					
	1 HOUR		2 HOURS		3 HOURS	
	light	dark	light	dark	light	dark
Empicol LX	0.1		0.6		1.0	
IPA	0.3		1.7		4.4	

Table 7

(Example 9)			
Ethanol %	Imbentin C91-35 %	Log (reduction)	
		After Light Exposure	No Rose Bengal
5	0.2	+6.8	
5	0.6	4.6	
10	0.6	+6.8	
15	0.6	+6.8	
5	-	+6.8	0.1
10	-	+6.8	-0.4
15	-	+6.8	0.0
-	0.2	4.6	2.5
-	0.6	3.5	2.3
-	-	+6.8	2.3

Table 8

(Example 10)			
Dowanol %	Imbentin C91-35 %	Log (reduction)	
		After Light Exposure	No Rose Bengal
3	0.7	+6.9	
3	-	+6.9	4.0
-	0.7	5.2	3.4
-	-	+6.9	

Table 9

(Example 11)			
Ethylene Glycol %	Imbentin C91-35 %	Log (reduction)	
		After Light Exposure	No Rose Bengal
10	0.7	+6.8	
10	-	+6.9	-0.2
-	0.7	+6.8	3.4
-	-	+6.8	

Table 10

(Example 12)			
Propan-2-ol %	Lialet 111 %	Log (reduction)	
		After Light Exposure	No Rose Bengal
15	0.5	+6.7	
15	-	+6.7	4.9
-	0.5	+6.7	+6.7
-	-	+6.7	

Table 11

(Example 13)			
Propan-2-ol %	Imbentin C91-35 %	Log (reduction)	
		After Light Exposure	No Rose Bengal
5	0.1	2.3	
10	0.1	+6.8	
10	0.5	+6.8	
10	0.7	+6.8	
-	-	4.8	
5	-	5.5	0.3
10	-	3.0	1.9
-	0.1	1.2	1.3
-	0.5	1.0	1.2
-	0.7	1.1	1.1

Table 12

(Example 14)			
Ethanol %	Imbentin C91-35 %	Log (reduction)	
		After Light Exposure	No Rose Bengal
5	0.2	4.1	
15	0.6	+7.1	
5	-	5.0	0.2
10	-	+7.1	0.2
15	-	+7.1	2.2
-	0.2	3.3	2.5
-	0.6	3.4	2.6
-	-	3.9	

Claims

1. A method of killing bacteria in biofilms on a surface, comprising applying to the surface a composition comprising a dyestuff which is capable of photo-dynamic inactivation of micro-organisms.
2. A method according to claim 1, wherein the dyestuff generates singlet oxygen on exposure to light.
3. A method according to claim 1 or 2, wherein the dyestuff is substantive to micro-organisms.
4. A method according to claim 1, 2 or 3, wherein the dyestuff is bleached by exposure to light.
5. A method according to any one of the preceding claims, wherein the dyestuff is selected from the group comprising Rose Bengal, Erythrosin B and phthalocyanin sulphonates.
6. A method according to any one of the preceding claims, wherein dyestuff is present in an amount in the range 1 to 100 ppm.
7. A method according to any one of the preceding claims, wherein the composition further comprises one or more surfactants.
8. A method according to claim 7, wherein the surfactant is alkoxyated.
9. A method according to claim 8, wherein the surfactant is ethoxyated.
10. A method according to claim 7, 8 or 9, wherein the surfactant is at least predominantly non-ionic and/or anionic.
11. A method according to any one of claims 7 to 10, wherein surfactant is present in the composition in an amount in the range 0.05 to 2.5% by weight of the total weight of the composition.
12. A method according to any one of the preceding claims, wherein the composition comprises one or more solvents.
13. A method according to claim 12, wherein the solvent is polar.
14. A method according to claim 13, wherein the solvent is a straight or branched chain C2 to C5 alcohol.
15. A method according to claim 12, 13 or 14, wherein solvent is present in the composition in an amount in the range 2 to 20% by weight of the total weight of the composition.

16. A method according to any one of the preceding claims, wherein the composition has a pH in the range 3 to 5.
17. A method according to claim 16, wherein the composition has a pH of about 4.
- 5 18. A method of cleaning a surface and killing bacteria in biofilms on the surface, comprising applying to the surface a composition comprising a dyestuff which is capable of photo-dynamic inactivation of micro-organisms, a surfactant and a solvent.

Patentansprüche

- 10 1. Verfahren zur Abtötung von Bakterien in biologischen Filmen auf einer Oberfläche umfassend, daß man auf die Oberfläche eine Zusammensetzung aufträgt, die einen Farbstoff enthält, der Mikroorganismen photodynamisch inaktivieren kann.
- 15 2. Verfahren nach Anspruch 1, worin der Farbstoff Singulett-Sauerstoff bei Belichtung erzeugt.
3. Verfahren nach Anspruch 1 oder Anspruch 2, worin der Farbstoff auf Mikroorganismen substantiv ist.
- 20 4. Verfahren nach einem der Ansprüche 1, 2 oder 3, worin der Farbstoff durch Belichtung gebleicht wird.
5. Verfahren nach einem der vorhergehenden Ansprüche, worin der Farbstoff ausgewählt ist aus der Gruppe umfassend Bengalrosa, Erythrosin B und Phthalocyaninsulfonate.
- 25 6. Verfahren nach einem der vorhergehenden Ansprüche, worin der Farbstoff in einer Menge im Bereich von 1 bis 100 ppm vorhanden ist.
7. Verfahren nach einem der vorhergehenden Ansprüche, worin die Zusammensetzung weiterhin ein oder mehrere Tenside umfaßt.
- 30 8. Verfahren nach Anspruch 7, worin das Tensid alkoxyliert ist.
9. Verfahren nach Anspruch 8, worin das Tensid ethoxyliert ist.
- 35 10. Verfahren nach Anspruch 7, 8 oder 9, worin das Tensid mindestens überwiegend nichtionisch und/oder anionisch ist.
11. Verfahren nach einem der Ansprüche 7 bis 10, worin das Tensid in der Zusammensetzung in einer Menge im Bereich von 0,05 bis 2,5 Gew.-%, bezogen auf die Gesamtmenge der Zusammensetzung, vorhanden ist.
- 40 12. Verfahren nach einem der vorhergehenden Ansprüche, worin die Zusammensetzung ein oder mehrere Lösungsmittel umfaßt.
13. Verfahren nach Anspruch 12, worin das Lösungsmittel polar ist.
- 45 14. Verfahren nach Anspruch 13, worin das Lösungsmittel ein geradkettiger oder verzweigter C₂-C₅-Alkohol ist.
15. Verfahren nach einem der Ansprüche 12, 13 oder 14, worin das Lösungsmittel in der Zusammensetzung in einer Menge im Bereich von 2 bis 20 Gew.-%, bezogen auf das Gesamtgewicht der Zusammensetzung, vorhanden ist.
- 50 16. Verfahren nach einem der vorhergehenden Ansprüche, worin die Zusammensetzung einen pH im Bereich von 3 bis 5 hat.
17. Verfahren nach Anspruch 16, worin die Zusammensetzung einen pH von etwa 4 hat.
- 55 18. Verfahren zum Reinigen einer Oberfläche und zum Abtöten von Bakterien in biologischen Filmen auf einer Oberfläche, umfassend, daß man auf die Oberfläche eine Zusammensetzung aufträgt, die einen Farbstoff, der Mikroorganismen photodynamisch inaktivieren kann, ein Tensid und ein Lösungsmittel enthält.

Revendications

1. Un procédé de destruction des bactéries dans des biofilms sur une surface, comprenant l'application sur la surface d'une composition contenant une substance colorante qui est capable d'inactiver des micro-organismes par voie photodynamique.
2. Un procédé selon la revendication 1, dans lequel la substance colorante génère de l'oxygène singulet à l'exposition à la lumière.
3. Un procédé selon la revendication 1 ou 2, dans lequel la substance colorante est substantive aux micro-organismes.
4. Un procédé selon la revendication 1, 2 ou 3, dans lequel la substance colorante est blanchie par exposition à la lumière.
5. Un procédé selon l'une quelconque des revendications précédentes, dans lequel la substance colorante est sélectionnée parmi le groupe composé de rose Bengale, d'érythrosine B et de sulfonates de phtalocyanine.
6. Un procédé selon l'une quelconque des revendications précédentes, dans lequel la substance colorante est présente dans une quantité comprise dans la gamme allant de 1 à 100 ppm.
7. Un procédé selon l'une quelconque des revendications précédentes, dans lequel la composition comprend en outre un ou plusieurs agent(s) tensioactif(s).
8. Un procédé selon la revendication 7, dans lequel l'agent tensioactif est alcoxylé.
9. Un procédé selon la revendication 8, dans lequel l'agent tensioactif est éthoxylé.
10. Un procédé selon la revendication 7, 8 ou 9, dans lequel l'agent tensioactif est au moins majoritairement non ionique et/ou anionique.
11. Un procédé selon l'une quelconque des revendications 7 à 10, dans lequel l'agent tensioactif est présent dans la composition dans une quantité comprise dans la gamme allant de 0,05 à 2,5 % en masse de la masse totale de la composition.
12. Un procédé selon l'une quelconque des revendications précédentes, dans lequel la composition comprend un ou plusieurs solvant(s).
13. Un procédé selon la revendication 12, dans lequel le solvant est polaire.
14. Un procédé selon la revendication 13, dans lequel le solvant est un alcool en C₂-C₅ à chaîne linéaire ou ramifiée.
15. Un procédé selon la revendication 12, 13 ou 14, dans lequel le solvant est présent dans la composition dans une quantité comprise dans la gamme allant de 2 à 20 % en masse de la masse totale de la composition.
16. Un procédé selon l'une quelconque des revendications précédentes, dans lequel la composition a un pH compris dans la gamme allant de 3 à 5.
17. Un procédé selon la revendication 16, dans lequel la composition a un pH de l'ordre de 4.
18. Un procédé de nettoyage d'une surface et de destruction des bactéries dans des biofilms sur la surface, comprenant l'application sur la surface d'une composition comprenant une substance colorante qui est capable d'inactiver par voie photodynamique des micro-organismes, un agent tensioactif et un solvant.




Fig. 1a




Fig. 1b





Fig 2a



* CONTROL (G-) * E.coli ▽ CONTROL (G+) ▲ S.aureus

Fig. 2b




Fig. 3a




Fig. 3b




Fig. 4a




Fig. 4b




Fig. 5




Fig. 6a




Fig. 6b




Fig. 7a




Fig. 7b




Fig. 8




Fig. 9




Fig. 10